

MICHAEL STEWART, MD, FRCPC



Dr. Michael Stewart is an Assistant Professor of Medicine at Dalhousie University and a consultant gastroenterologist at the QEII Health Sciences Centre in Halifax, Nova Scotia. Dr. Stewart completed his undergraduate medical education at Dalhousie University, residency at the University of Calgary and Dalhousie University, and a fellowship in Inflammatory Bowel Disease at Cedars-Sinai Medical Centre. His clinical and research interests include intestinal ultrasound, colon cancer, system innovation, and clinical trials. He is the co-Director of the Nova Scotia Collaborative IBD Program and Medical Director of the Nova Scotia Colon Cancer Prevention Program.

Affiliations: Assistant Professor, Division of Digestive Care & Endoscopy, Department of Medicine, Dalhousie University, Halifax, NS

TREATMENT RELATED ADVERSE EVENTS AND MONITORING OF PATIENTS RECEIVING BIOLOGIC OR SMALL MOLECULE THERAPY FOR INFLAMMATORY BOWEL DISEASE.

Key Takeaways

- Advanced therapies for IBD are generally safe but require treatment specific ongoing monitoring.
- Individual patient characteristics influence treatment choice and should be considered when implementing an ongoing monitoring strategy.
- Regular biochemical monitoring should be individualized to specific treatment requirements.
- Drug interactions must be considered when prescribing small molecule advanced therapy for IBD.

Introduction

The management of Inflammatory Bowel Disease (IBD) has evolved with the emergence of new treatment paradigms and the introduction of novel advanced therapies, including monoclonal antibodies (mAbs) and small molecules. These advanced therapies have improved disease control, but they necessitate careful pre-treatment assessment and ongoing monitoring to manage potential adverse effects and optimize patient outcomes. This review focuses on practical approaches to treatment-specific monitoring of currently available advanced therapies.

Treatment-associated adverse events

Infections

Patients with IBD, and those taking advanced therapies, are at an increased risk for infections. Maintaining vigilance for signs of infection, prompt evaluation and management, and therapy interruption, when necessary, are crucial in avoiding serious complications.

The risk of opportunistic infections is a significant concern with anti-tumor necrosis factor (TNF) therapy, because this treatment method doubles the risk of such infections among IBD patients.¹ In addition, the risk of tuberculosis reactivation can increase up to 25-fold depending on clinical circumstances.² The TREAT Registry showed a serious infection rate of 2.15 events per 100 patient-years (PY).³ However, a meta-analysis of 21 placebo-controlled Crohn's disease (CD) trials did not show an increased risk of serious infections with anti-TNF therapy.⁴ Excluding latent infections prior to treatment and ongoing monitoring, especially for opportunistic and atypical infections, is important when administering anti-TNF therapy.

In a phase 3 CD trial, nasopharyngitis was more frequent in the vedolizumab arm, along with apparently higher rates of both infections and serious infections.⁵ However, subsequent long-term safety studies and meta-analyses did not show an increased infection risk with vedolizumab.⁶⁻⁹ The EVOLVE study, a multicenter retrospective real-world study that included 1,095 IBD patients, found a significantly lower rate of serious infections and adverse events with vedolizumab versus anti-TNF.¹⁰

Ustekinumab therapy has not shown an increased risk for serious or opportunistic infections in long-term studies,¹¹ with its infection risk being similar to vedolizumab and lower than that of anti-TNF therapies.^{12,13} Risankizumab and mirikizumab therapy have also shown no increased risk of serious or opportunistic infections in the registrational clinical trials.¹⁴⁻¹⁶

The introduction of Janus kinase (JAK) inhibitors has raised specific concerns around Herpes Zoster (HZ) reactivation. Long-term data on tofacitinib suggests that HZ occurs at a rate of 3.²⁴ events per 100 PY, with other serious infections occurring at a rate of 1.⁸ events per 100 PY.¹⁷ An upadacitinib trial reported similar serious infection rates to adalimumab, but a higher risk of HZ.¹⁸ A recent network meta-analysis concluded that tofacitinib and upadacitinib significantly increase the risk of HZ infection,¹⁹ although most cases were reported to be mild or moderate and had resolved without discontinuation of treatment.²⁰ Routine use of the adjuvanted recombinant zoster vaccine is recommended for adults requiring advanced IBD therapies.

In a phase 3 trial for UC, it was observed that ozanimod exhibited infection rates of 23% (compared to 11.9% with placebo), with low rates of serious infections (0.9% for ozanimod versus 1.8% with placebo) and HZ (2.2% for ozanimod versus 0.4% with placebo).²¹ These results were confirmed by a long-term extension study that reported an infection rate of 24.³ events per 100 PY, a serious infection rate of 1.9 events per 100 PY, and an HZ rate of 1.7 events per 100 PY.²² Notably, an open-label study involving multiple sclerosis patients highlighted that opportunistic infections were predominantly driven by HZ.²³ Similarly, in a phase 3 trial for UC, it was found that etrasimod demonstrated minimal serious infection rates (1% for etrasimod versus 3% for placebo) and HZ rates (1% for etrasimod versus 0% for placebo), with no reports of opportunistic infections.²⁴ Consistent with these findings, long-term safety data from an etrasimod open-label extension trial indicated a low risk of infection.²⁵

Vaccination Status

Live vaccines are contraindicated in patients receiving biologic and small molecule therapy. It is important to assess measles, mumps, and rubella (MMR) and varicella-zoster virus (VZV) vaccination history and immune status before initiating advanced therapy. If required, administer vaccines before starting therapy; however, do not delay urgent treatment for live vaccine administration.

Recommendations indicate that all IBD patients should receive the following inactivated vaccines, regardless of active treatment: influenza, meningococcal, *Haemophilus influenzae* type b, diphtheria, tetanus, pertussis, human papillomavirus, and pneumococcal.

Assess viral hepatitis status before initiating advanced therapy for IBD. Unimmunized patients should receive the hepatitis B vaccine. It is important to note that reactivation of hepatitis B is a known complication of anti-TNF therapy. Patients positive for hepatitis B surface antigen (HBsAg) are at the highest risk and should consider prophylactic antiviral therapy before initiating anti-TNF treatment.

All adult IBD patients should consider the recombinant zoster vaccine (non-live), especially those receiving immunomodulator, biologic, or small molecule therapy as it can mitigate HZ risk.

For further details on immunizations for IBD patients, refer to the 2021 Canadian Association of Gastroenterology Clinical Practice Guideline.²⁶

Hematologic And Metabolic

Up to 19% of patients receiving anti-TNF therapy for immune-mediated diseases develop at least one episode of neutropenia, with 6% experiencing serious infections related to neutropenia.²⁷ Thrombocytopenia is infrequently associated with anti-TNF therapies, with data limited to case reports. In cases of significant thrombocytopenia, alternate causes, including autoimmune conditions or viral infections, should be considered.²⁸

Weight gain has been observed in patients with IBD who are receiving anti-TNF therapy. However, long-term registry data has not established a direct link between anti-TNF therapy and weight gain, although patients who are underweight at treatment initiation may experience early weight gain.²⁹ Some patients gain weight due to an improvement in their nutritional status following effective therapy, as suggested by a small cohort study that showed an increase in both body mass index (BMI) and muscle mass parameters after anti-TNF therapy initiation.³⁰

Vedolizumab therapy for IBD has not been associated with metabolic adverse effects. While leukocytosis and leukopenia were reported in a small proportion of patients in registration trials, subsequent long-term safety analyses have not confirmed these findings.⁹ Therapies that target interleukins do not appear to cause significant adverse hematologic or metabolic effects.^{15,16,31,32}

Neutropenia and lymphopenia occurred in upadacitinib-treated patients in the pivotal induction and maintenance trials, with no cases requiring treatment discontinuation. Neutropenia was observed in 6% of patients treated with 30 mg of upadacitinib, 3% of patients treated with 15 mg of upadacitinib, and in 1% of patients who received a placebo. Lymphopenia occurred in 2% of patients who were treated with both 30 mg and 15 mg doses of upadacitinib, and in 1% of placebo-treated patients. Anemia was more common in placebo-treated patients compared to those receiving upadacitinib.³³ With up to 9.2 years of safety data, significant cytopenias have not been reported with tofacitinib.¹⁷ Creatine phosphokinase (CPK) elevations were observed in a small percentage of JAK inhibitor

patients and were mostly asymptomatic and non-serious.

S1P receptor modulators impair the migration of lymphocytes out of lymphoid tissue by blocking S1P receptors, leading to a relative reduction in circulating peripheral lymphocytes. There is generally an expected and measurable relative reduction in lymphocytes by approximately 40%–50%, which resolves after treatment discontinuation in most patients. Profound lymphopenia is rare, occurring in 1% of patients.^{21,25,34}

Cardiovascular

Patients with IBD are at an increased risk of cardiovascular disease,³⁵ likely attributable to chronic inflammation and associated metabolic derangement.³⁶ While effective management of IBD and its underlying risk factors is key, there are specific treatment-related considerations.

Data from preclinical studies suggested potential benefits of TNF α inhibition for treating congestive heart failure, however, a subsequent clinical trial showed no such benefit, and had reported an increased risk of hospitalization and all-cause mortality.³⁷ Case reports also link anti-TNF therapy to heart failure exacerbations in patients with IBD.³⁸ Anti-TNF therapy is contraindicated in New York Heart Association Class III/IV heart failure and should be used with caution in patients at risk for heart failure.

Long-term safety data has not established an increased cardiovascular event risk with vedolizumab therapy.⁹ Agents targeting IL-12 and -23 show a favourable safety profile with no significant increase in cardiovascular events compared to other therapies.³⁹

Initiation of JAK inhibitor therapy can modestly increase both low density lipoprotein (LDL) and high-density lipoprotein (HDL) cholesterol levels by approximately 20%, with the LDL-HDL ratio remaining stable.^{33,40–42} It is not clear if increased cholesterol levels results in atherosclerosis. Interestingly, there is some evidence that tofacitinib may positively impact macrophage cholesterol metabolism, which could potentially mitigate the risk of atherosclerosis.⁴³

In a long-term extension study of tofacitinib for treating UC, the risk of major adverse cardiovascular events was low, with a rate of 0.27 events per 100 PY.¹⁷ A systematic review and meta-analysis of real-world studies also did not report any major adverse cardiovascular events or thromboembolic complications.⁴⁴

The ORAL Surveillance open-label randomized trial compared tofacitinib at a dose of 5 mg or 10 mg twice daily to anti-TNF therapy in 4,362 patients older than 50 years with active rheumatoid arthritis and at least one additional cardiovascular risk factor. The results of the trial demonstrated a higher incidence of major cardiovascular events with tofacitinib.⁴⁵ Post-hoc analyses demonstrated that the increased risk of adverse cardiovascular events was limited to a high-risk patient cohort (age ≥ 65 years or those with a history of smoking)⁴⁶ and was predominantly observed

in patients with prior atherosclerotic cardiovascular disease.⁴⁷ The SELECT-COMPARE trial compared the effects of upadacitinib and adalimumab for rheumatoid arthritis treatment, and found no difference in the incidence of cardiovascular events.⁴⁸

S1PRMs pose specific cardiac safety concerns due to S1P1 receptors, which are found on cardiac myocytes, and their subsequent effects on cardiac conduction. Transient bradycardia is a common early side effect, within hours of the first dose, which is largely asymptomatic. In the True North induction and maintenance trials, one patient developed a type 1 second-degree heart block, and there were no cases of type II or third-degree heart block.^{21,34} A large open-label extension trial of ozanimod for multiple sclerosis reported hypertension at a rate of 2.0 events per 100 PY and no cases of second- or third-degree heart block.²³

Thromboembolic

IBD has long been recognized as a risk factor for venous thromboembolism (VTE) and arterial events, especially during disease exacerbation.^{49–51} Corticosteroid use increases VTE risk, while anti-TNF agents have been associated with a decreased risk of VTE.^{51,52}

Despite regulatory warnings prompted by the ORAL Surveillance study, long-term exposure data suggests that the risk of VTE and arterial thrombosis in those treated with JAK inhibitors remains low. Randomized trials and real-world studies have consistently found low rates of these adverse events that do not differ from those observed with anti-TNF therapy.^{17,44,48,53,54} A recent consensus process concluded that there is no observable increased risk of VTE in IBD patients treated with tofacitinib.⁵¹

Hepatic

Anti-TNF therapies have been associated with a variety of liver injury patterns, with events ranging from transient and self-limited, to severe.⁵⁵ Anti-integrin and anti-interleukin therapies have a low risk of drug-induced liver injury, although there have been cases of idiosyncratic, clinically apparent liver injury that has resolved with discontinuation.^{9,56}

Unlike monoclonal antibodies, small molecule drugs undergo hepatic metabolism through the cytochrome P450 enzyme system, which can result in drug-drug interactions. Elevations of transaminases have been observed with both JAKs and S1PRMs, although they are generally mild and do not require treatment discontinuation.^{22,24,33,56}

Neurologic

Anti-TNF agents increase the risk of inflammatory demyelinating and non-demyelinating central nervous system (CNS) events, especially in patients with multiple sclerosis or a history of optic neuritis.⁵⁷ Other advanced therapies do not appear to increase the risk of inflammatory CNS events.

One case of progressive multifocal leukoencephalopathy (PML) has been reported in a vedolizumab-treated patient who was HIV-positive and on concomitant immunosuppression, another case of PML was reported in an infliximab-treated patient,⁵⁸ and there have been case reports of PML in S1PRM-treated multiple sclerosis patients.²³

Posterior reversible encephalopathy syndrome (PRES) has been reported in patients treated with anti-TNF agents^{59,60} and ustekinumab.^{61,62}

Ophthalmologic

Clinical trial data suggest that S1PRMs can trigger macular edema in 1:125 to 1:300 patients.^{21,22,34} The cases generally resolve following drug discontinuation, however, patients with pre-existing uveitis or diabetes are at increased risk.

Immunologic

Anti-TNF therapy triggers a spectrum of immune-mediated adverse events, including infusion reactions, injection site reactions, delayed hypersensitivity reactions, paradoxical autoimmune disorders (e.g., lupus-like syndromes and psoriasis), and immunogenicity. Subsequent mAbs and small molecule

therapies have largely attenuated these immunologic complications of treatment.

Malignancy

The use of anti-TNF agents has raised concerns around an increased risk of malignancy, specifically non-melanoma skin cancer (NMSC) and lymphoma,⁶³ although the evidence has been conflicting.^{64,65} The S1PRM modulator fingolimod has a slightly increased risk of basal cell carcinoma,⁶⁶ which has not been conclusively demonstrated with ozanimod or etrasimod.^{21,22,25} Findings on the malignancy risk of JAKs are also varied, with some studies suggesting a risk of malignancy and NMSC.^{17,40,67,68}

Treatment Monitoring Strategy

Effective IBD management requires a baseline assessment and ongoing monitoring for treatment-related complications. Regular laboratory investigations, symptom monitoring, infection vigilance, cancer screening, and attention to treatment-specific concerns are crucial. Please see the table below for more information.

Therapeutic Class	Medication	Pre-Treatment Assessment	Ongoing Monitoring
Anti-tumour necrosis factor-α (TNFα)	Infliximab	CBC, hepatic function, viral hepatitis (HBV, HCV), TB status, exposure to opportunistic pathogens. Vaccine review (no live vaccination during treatment). Contraindicated if: - Active infection. - Profound cytopenia. - NYHA Class III or IV heart failure. - Pre-existing multiple sclerosis or optic neuritis.	CBC every 3–6 months. Liver panel every 3–6 months. Tb/viral hepatitis if high-risk travel or exposure. Monitor for signs and symptoms of infection with consideration of atypical/opportunistic pathogens. Consider an annual pap-smear and skin exam, especially if concomitant immunosuppressive therapy. Annual influenza vaccine and COVID-19 vaccine as per National Advisory Committee on Immunization (NACI) recommendations.
	Adalimumab		
	Golimumab		
	Certolizumab		
Anti-integrin	Vedolizumab	Vaccine review (no live vaccination during treatment). Consider TB status assessment.	CBC every 3–6 months. Liver panel every 3–6 months. Annual influenza vaccine and COVID-19 vaccine as per National Advisory Committee on Immunization (NACI) recommendations.
Anti-interleukin	Ustekinumab	CBC, hepatic function, viral hepatitis (HBV, HCV), TB status, exposure to opportunistic pathogens. Vaccine review (no live vaccination during treatment).	CBC every 3–6 months. Liver panel every 3–6 months. Monitor for signs and symptoms of infection with consideration of atypical/opportunistic pathogens. Annual influenza vaccine and COVID-19 vaccine as per National Advisory Committee on Immunization (NACI) recommendations.
	Risankizumab		
	Mirikizumab		

Janus Kinase inhibitor	Tofacitinib	CBC, hepatic function, viral hepatitis (HBV, HCV), TB status, exposure to opportunistic pathogens. Baseline lipid panel and cardiovascular risk factor assessment. If age >65 years or history of cardiovascular disease, use lowest effective dose with careful consideration of risks and benefits.	CBC every 3–6 months: - Interrupt treatment if HGB <80 g/L or decrease >20 g/L; or ANC $0.5\text{--}1.0 \times 10^9$ cells/L. - Discontinue if ANC <0.5×10^9 cells/L or ALC <0.5×10^9 cells/L. - Liver panel at 4–8 weeks, then every 3–6 months. Lipid panel at week 4–8 (tofacitinib)/week 12 (upadacitinib); then every 6 months.
	Upadacitinib	Family planning, if applicable. - Dose adjustment (5 mg BID) if: - eGFR <60. - Strong CYP3A4 inhibitors. - Moderate CYP3A4 inhibitor with a strong CYP2C19 inhibitor. Contraindicated if: - Pre-existing cytopenia (ANC <1.0×10^9 cells/L, HGB <90 g/L, ALC <0.5×10^9 cells/L). - Severe renal (eGFR <15 ml/min) or hepatic impairment. - Potent CYP3A4 inducers. Vaccine review (recombinant herpes zoster highly recommended, no live vaccines during treatment).	Coordinate hypercholesterolemia management with primary care/ cardiology, per 2021 Canadian Cardiovascular Society Guidelines. Periodic confirmation of medication adherence. Periodic review of family planning, if applicable. Monitor for signs and symptoms of infection with consideration of atypical/opportunistic pathogens. Consider an annual skin exam. Annual influenza vaccine and COVID-19 vaccine as per National Advisory Committee on Immunization (NACI) recommendations.
Sphingosine-1-phosphate receptor (S1PR) modulators	Ozanimod	CBC, hepatic function, viral hepatitis (HBV, HCV), TB status, exposure to opportunistic pathogens. Cardiac assessment: ECG, heart rate, blood pressure. Ophthalmology evaluation (if diabetes mellitus, uveitis, or retinal disease). Family planning, if applicable. Caution if: - Pre-existing pulmonary disease. - Drugs that slow the heart rate or AV conduction.	CBC every 3–6 months: Interrupt treatment if ALC < 0.2×10^9 cells/L Liver panel every 3–6 months. Assess visual disturbances. Monitor blood pressure regularly. Periodic confirmation of medication adherence.
	Etrasimod	Contraindicated if: - Concomitant use of MAO inhibitors. - Severe hepatic impairment. - Myocardial infarction, unstable angina, stroke, or transient ischemic attack, decompensated or advanced heart failure, within 6 months. - Cardiac conduction abnormalities (AV node block, SA block) without a pacemaker. - Macular edema. - Severe respiratory disease (pulmonary fibrosis, asthma, or chronic obstructive pulmonary disease); spirometry if indicated. Vaccine review (recombinant Herpes Zoster highly recommended, no live vaccines during treatment).	Periodic review of family planning, if applicable. Monitor for signs and symptoms of infection with consideration of atypical/opportunistic pathogens. Consider annual skin exam. Annual influenza vaccine and COVID-19 vaccine as per National Advisory Committee on Immunization (NACI) recommendations

ALC, absolute lymphocyte count; ANC, absolute neutrophil count; AV, atrioventricular; BID, twice a day; CBC, complete blood count; CYP, cytochrome P450; eGFR, estimated glomerular filtration rate; ECG, electrocardiogram; HBV, Hepatitis B virus; HCV, Hepatitis C virus; HGB, hemoglobin; NYHA, New York Heart Association; SA, sinoatrial; Tb, tuberculosis; TNF α , tumour necrosis factor- α .

Table 1: Therapeutic class-based guide for advance therapy monitoring in the management of Inflammatory Bowel Disease ; courtesy of Michael Stewart, MD, FRCPC

Correspondence:

Michael Stewart, MD, FRCPC
Email: Michael.stewart@dal.ca

Financial Disclosures:

Grants/Research Support: Abbvie, Janssen, Takeda
Speakers Bureau/Honoraria: Abbvie, Takeda, Janssen, Eli Lilly
Consulting Fees: Abbvie, Takeda, Janssen, Pfizer, Sandoz, Bristol-Myer-Squibb, Eli Lilly, Celltrion

References:

1. Ford AC, Peyrin-Biroulet L. Opportunistic infections with anti-tumor necrosis factor- α therapy in inflammatory bowel disease: meta-analysis of randomized controlled trials. *Am J Gastroenterol*. 2013;108(8):1268-1276. doi:10.1038/ajg.2013.138
2. Solovic I, Sester M, Gomez-Reino JJ, Rieder HL, Ehlers S, Milburn HJ, et al. The risk of tuberculosis related to tumour necrosis factor antagonist therapies: a TBNET consensus statement. *European Respiratory Journal*. 2010;36(5):1185-1206. doi:10.1183/09031936.00028510
3. Lichtenstein GR, Feagan BG, Cohen RD, Salzberg BA, Safdi M, Popp JW, Jr., et al. Infliximab for Crohn's disease: more than 13 years of real-world experience. *Inflamm Bowel Dis*. 2018;24(3):490-501. doi:10.1093/ibd/izx072
4. Peyrin-Biroulet L, Deltenre P, de Suray N, Branche J, Sandborn WJ, Colombel JF. Efficacy and safety of tumor necrosis factor antagonists in Crohn's disease: meta-analysis of placebo-controlled trials. *Clin Gastroenterol Hepatol*. 2008;6(6):644-653. doi:10.1016/j.cgh.2008.03.014
5. Sandborn WJ, Feagan BG, Rutgeerts P, Hanauer S, Colombel JF, Sands BE, et al. Vedolizumab as induction and maintenance therapy for Crohn's disease. *N Engl J Med*. 2013;369(8):711-721. doi:10.1056/NEJMoa1215739
6. Shah ED, Farida JP, Siegel CA, Chong K, Melmed GY. Risk for overall infection with anti-TNF and anti-integrin agents used in IBD: a systematic review and meta-analysis. *Inflamm Bowel Dis*. 2017;23(4):570-577. doi:10.1097/mib.0000000000001049
7. Luthra P, Peyrin-Biroulet L, Ford AC. Systematic review and meta-analysis: opportunistic infections and malignancies during treatment with anti-integrin antibodies in inflammatory bowel disease. *Aliment Pharmacol Ther*. 2015;41(12):1227-1236. doi:10.1111/apt.13215
8. Colombel JF, Sands BE, Rutgeerts P, Sandborn W, Danese S, D'Haens G, et al. The safety of vedolizumab for ulcerative colitis and Crohn's disease. *Gut*. 2017;66(5):839-851. doi:10.1136/gutjnl-2015-311079
9. Loftus EV, Jr., Feagan BG, Panaccione R, Colombel JF, Sandborn WJ, Sands BE, et al. Long-term safety of vedolizumab for inflammatory bowel disease. *Aliment Pharmacol Ther*. 2020;52(8):1353-1365. doi:10.1111/apt.16060
10. Bressler B, Yarur A, Silverberg MS, Bassel M, Bellaguarda E, Fourment C, et al. Vedolizumab and anti-tumour necrosis factor α real-world outcomes in biologic-naïve inflammatory bowel disease patients: results from the EVOLVE study. *J Crohns Colitis*. 2021;15(10):1694-1706. doi:10.1093/ecco-jcc/jjab058
11. Sandborn WJ, Rebuck R, Wang Y, Zou B, Adedokun OJ, Gasink C, et al. Five-year efficacy and safety of ustekinumab treatment in Crohn's disease: the IM-UNITI trial. *Clin Gastroenterol Hepatol*. 2022;20(3):578-590.e574. doi:10.1016/j.cgh.2021.02.025
12. Gebeyehu GG, Fiske J, Liu E, Limdi JK, Broglio G, Selinger C, et al. Ustekinumab and vedolizumab are equally safe and effective in elderly Crohn's disease patients. *Dig Dis Sci*. 2023;68(5):1983-1994. doi:10.1007/s10620-022-07770-8
13. Cheng D, Kochar BD, Cai T, Ananthakrishnan AN. Risk of infections with ustekinumab and tofacitinib compared to tumor necrosis factor α antagonists in inflammatory bowel diseases. *Clin Gastroenterol Hepatol*. 2022;20(10):2366-2372.e2366. doi:10.1016/j.cgh.2022.01.013
14. D'Haens G, Panaccione R, Baert F, Bossuyt P, Colombel JF, Danese S, et al. Risankizumab as induction therapy for Crohn's disease: results from the phase 3 ADVANCE and MOTIVATE induction trials. *Lancet*. 2022;399(10340):2015-2030. doi:10.1016/s0140-6736(22)00467-6
15. Ferrante M, Panaccione R, Baert F, Bossuyt P, Colombel JF, Danese S, et al. Risankizumab as maintenance therapy for moderately to severely active Crohn's disease: results from the multicentre, randomised, double-blind, placebo-controlled, withdrawal phase 3 FORTIFY maintenance trial. *Lancet*. 2022;399(10340):2031-2046. doi:10.1016/s0140-6736(22)00466-4
16. D'Haens G, Dubinsky M, Kobayashi T, Irving PM, Howaldt S, Pokrotnieks J, et al. Mirikizumab as induction and maintenance therapy for ulcerative colitis. *N Engl J Med*. 2023;388(26):2444-2455. doi:10.1056/NEJMoa2207940
17. Panés J, D'Haens GR, Sands BE, Ng SC, Lawendy N, Kulisek N, et al. Analysis of tofacitinib safety in ulcerative colitis from the completed global clinical developmental program up to 9.2 years of drug exposure. *United European Gastroenterol J*. 2024. doi:10.1002/ueg.212584
18. Burmester GR, Cohen SB, Winthrop KL, Nash P, Irvine AD, Deodhar A, et al. Safety profile of upadacitinib over 15 000 patient-years across rheumatoid arthritis, psoriatic arthritis, ankylosing spondylitis and atopic dermatitis. *RMD Open*. 2023;9(1). doi:10.1136/rmdopen-2022-002735
19. Din S, Selinger CP, Black CJ, Ford AC. Systematic review with network meta-analysis: risk of Herpes Zoster with biological therapies and small molecules in inflammatory bowel disease. *Aliment Pharmacol Ther*. 2023;57(6):666-675. doi:10.1111/apt.17379
20. Winthrop KL, Vermeire S, Long MD, Panés J, Ng SC, Kulisek N, et al. Long-term risk of Herpes Zoster infection in patients with ulcerative colitis receiving tofacitinib. *Inflamm Bowel Dis*. 2023;29(1):85-96. doi:10.1093/ibd/izac063
21. Sandborn WJ, Feagan BG, D'Haens G, Wolf DC, Jovanovic I, Hanauer SB, et al. Ozanimod as induction and maintenance therapy for ulcerative colitis. *N Engl J Med*. 2021;385(14):1280-1291. doi:10.1056/NEJMoa2033617
22. Danese S, Panaccione R, Abreu MT, Rubin DT, Ghosh S, Dignass A, et al. Efficacy and safety of approximately 3 years of continuous ozanimod in moderately to severely active ulcerative colitis: interim analysis of the True North open-label extension. *J Crohns Colitis*. 2024;18(2):264-274. doi:10.1093/ecco-jcc/jjad146
23. Cree BA, Selmaj KW, Steinman L, Comi G, Bar-Or A, Arnold DL, et al. Long-term safety and efficacy of ozanimod in relapsing multiple sclerosis: Up to 5 years of follow-up in the DAYBREAK open-label extension trial. *Mult Scler*. 2022;28(12):1944-1962. doi:10.1177/13524585221102584
24. Regueiro M, Siegmund B, Yarur AJ, Steinwurz F, Gecse KB, Goetsch M, et al. Etrasimod for the treatment of ulcerative colitis: analysis of infection events from the ELEVATE UC Clinical Program. *J Crohns Colitis*. 2024. doi:10.1093/ecco-jcc/jjae060
25. Vermeire S, Chiorean M, Panés J, Peyrin-Biroulet L, Zhang J, Sands BE, et al. Long-term safety and efficacy of etrasimod for ulcerative colitis: results from the open-label extension of the OASIS Study. *J Crohns Colitis*. 2021;15(6):950-959. doi:10.1093/ecco-jcc/jjab016
26. Jones JL, Tse F, Carroll MW, deBruyn JC, McNeil SA, Pham-Huy A, et al. Canadian Association of Gastroenterology Clinical Practice Guideline for Immunizations in Patients With Inflammatory Bowel Disease (IBD)-Part 2: Inactivated Vaccines. *J Can Assoc Gastroenterol*. 2021;4(4):e72-e91.

doi:10.1093/jcag/gwab016

27. Hastings R, Ding T, Butt S, Gadsby K, Zhang W, Moots RJ, et al. Neutropenia in patients receiving anti-tumor necrosis factor therapy. *Arthritis Care Res (Hoboken)*. 2010;62(6):764-769. doi:10.1002/acr.20037
28. Bessissow T, Renard M, Hoffman I, Vermeire S, Rutgeerts P, Van Assche G. Review article: non-malignant haematological complications of anti-tumour necrosis factor alpha therapy. *Aliment Pharmacol Ther*. 2012;36(4):312-323. doi:10.1111/j.1365-2036.2012.05189.x
29. Winter RW, Friedman S, Nielsen J, Kjeldsen J, Nørgård BM, Larsen MD. Infliximab is not associated with a general long-term weight gain in patients with inflammatory bowel disease: a nationwide study. *Am J Gastroenterol*. 2022;117(5):777-784. doi:10.14309/ajg.0000000000001721
30. Csontos Á A, Molnár A, Piri Z, Katona B, Dakó S, Pálfi E, et al. The effect of anti-TNF α induction therapy on the nutritional status and dietary intake in inflammatory bowel disease. *J Gastrointest Liver Dis*. 2016;25(1):49-56. doi:10.15403/jgld.2014.1121.251.tnf
31. Davies SC, Nguyen TM, Parker CE, MacDonald JK, Jairath V, Khanna R. Anti-IL-12/23p40 antibodies for maintenance of remission in Crohn's disease. *Cochrane Database Syst Rev*. 2019;12(12):Cd012804. doi:10.1002/14651858.CD012804.pub2
32. Hanauer SB, Sandborn WJ, Feagan BG, Gasink C, Jacobstein D, Zou B, et al. IM-UNITI: three-year efficacy, safety, and immunogenicity of ustekinumab treatment of Crohn's Disease. *J Crohns Colitis*. 2020;14(1):23-32. doi:10.1093/ecco-jcc/ijz110
33. Danese S, Vermeire S, Zhou W, Pangan AL, Siffledeen J, Greenbloom S, et al. Upadacitinib as induction and maintenance therapy for moderately to severely active ulcerative colitis: results from three phase 3, multicentre, double-blind, placebo-controlled, randomised trials. *Lancet*. 2022;399(10341):2113-2128. doi:10.1016/s0140-6736(22)00581-5
34. Sandborn WJ, Vermeire S, Peyrin-Biroulet L, Dubinsky MC, Panes J, Yarur A, et al. Etrasimod as induction and maintenance therapy for ulcerative colitis (ELEVATE): two randomised, double-blind, placebo-controlled, phase 3 studies. *Lancet*. 2023;401(10383):1159-1171. doi:10.1016/s0140-6736(23)00061-2
35. Panhwar MS, Mansoor E, Al-Kindi SG, Sinh P, Katz J, Oliveira GH, et al. Risk of myocardial infarction in inflammatory bowel disease: a population-based National study. *Inflamm Bowel Dis*. 2019;25(6):1080-1087. doi:10.1093/ibd/izy354
36. Hansson GK. Inflammation, atherosclerosis, and coronary artery disease. *N Engl J Med*. 2005;352(16):1685-1695. doi:10.1056/NEJMc043430
37. Chung ES, Packer M, Lo KH, Fasanmade AA, Willerson JT. Randomized, double-blind, placebo-controlled, pilot trial of infliximab, a chimeric monoclonal antibody to tumor necrosis factor- α , in patients with moderate-to-severe heart failure: results of the anti-TNF Therapy Against Congestive Heart Failure (ATTACH) trial. *Circulation*. 2003;107(25):3133-3140. doi:10.1161/01.Cir.0000077913.60364.D2
38. Kwon HJ, Coté TR, Cuffe MS, Kramer JM, Braun MM. Case reports of heart failure after therapy with a tumor necrosis factor antagonist. *Ann Intern Med*. 2003;138(10):807-811. doi:10.7326/0003-4819-138-10-200305200-00008
39. Vuyyuru SK, Solitano V, Hogan M, MacDonald JK, Zayadi A, Parker CE, et al. Efficacy and safety of IL-12/23 and IL-23 inhibitors for Crohn's disease: systematic review and meta-analysis. *Dig Dis Sci*. 2023;68(9):3702-3713. doi:10.1007/s10620-023-08014-z
40. Sandborn WJ, Su C, Panes J. Tofacitinib as induction and maintenance therapy for ulcerative colitis. *N Engl J Med*. 2017;377(5):496-497. doi:10.1056/NEJMc1707500
41. Sleutjes JAM, Roeters van Lennep JE, van der Woude CJ, de Vries AC. Lipid changes after induction therapy in patients with inflammatory bowel disease: effect of different drug classes and inflammation. *Inflamm Bowel Dis*. 2023;29(4):531-538. doi:10.1093/ibd/izac100
42. Sandborn WJ, Lawendy N, Danese S, Su C, Loftus EV, Jr., Hart A, et al. Safety and efficacy of tofacitinib for treatment of ulcerative colitis: final analysis of OCTAVE Open, an open-label, long-term extension study with up to 7.0 years of treatment. *Aliment Pharmacol Ther*. 2022;55(4):464-478. doi:10.1111/apt.16712
43. Adorni MP, Papotti B, Borghi MO, Raschi E, Zimetti F, Bernini F, et al. Effect of the JAK/STAT inhibitor tofacitinib on macrophage cholesterol metabolism. *Int J Mol Sci*. 2023;24(16). doi:10.3390/ijms241612571
44. Taxonera C, Olivares D, Alba C. Real-world effectiveness and safety of tofacitinib in patients with ulcerative colitis: systematic review with meta-analysis. *Inflamm Bowel Dis*. 2022;28(1):32-40. doi:10.1093/ibd/izab011
45. Ytterberg SR, Bhatt DL, Mikuls TR, Koch GG, Fleischmann R, Rivas JL, et al. Cardiovascular and cancer risk with tofacitinib in rheumatoid arthritis. *N Engl J Med*. 2022;386(4):316-326. doi:10.1056/NEJMoa2109927
46. Kristensen LE, Danese S, Yndestad A, Wang C, Nagy E, Modesto I, et al. Identification of two tofacitinib subpopulations with different relative risk versus TNF inhibitors: an analysis of the open label, randomised controlled study ORAL Surveillance. *Ann Rheum Dis*. 2023;82(7):901-910. doi:10.1136/ard-2022-223715
47. Charles-Schoeman C, Buch MH, Dougados M, Bhatt DL, Giles JT, Ytterberg SR, et al. Risk of major adverse cardiovascular events with tofacitinib versus tumour necrosis factor inhibitors in patients with rheumatoid arthritis with or without a history of atherosclerotic cardiovascular disease: a post hoc analysis from ORAL Surveillance. *Ann Rheum Dis*. 2023;82(1):119-129. doi:10.1136/ard-2022-222259
48. Fleischmann R, Mysler E, Bessette L, Peterfy CG, Durez P, Tanaka Y, et al. Long-term safety and efficacy of upadacitinib or adalimumab in patients with rheumatoid arthritis: results through 3 years from the SELECT-COMPARE study. *RMD Open*. 2022;8(1). doi:10.1136/rmdopen-2021-002012
49. Grainge MJ, West J, Card TR. Venous thromboembolism during active disease and remission in inflammatory bowel disease: a cohort study. *Lancet*. 2010;375(9715):657-663. doi:10.1016/s0140-6736(09)61963-2
50. Bernstein CN, Nugent Z, Singh H. Persistently high rate of venous thromboembolic disease in inflammatory bowel disease: a population-based study. *Am J Gastroenterol*. 2021;116(7):1476-1484. doi:10.14309/ajg.0000000000001237
51. Olivera PA, Zuily S, Kotze PG, Regnault V, Al Awadhi S, Bossuyt P, et al. International consensus on the prevention of venous and arterial thrombotic events in patients with inflammatory bowel disease. *Nat Rev Gastroenterol Hepatol*. 2021;18(12):857-873. doi:10.1038/s41575-021-00492-8
52. Nguyen GC, Elnahas A, Jackson TD. The impact of preoperative steroid use on short-term outcomes following surgery for inflammatory bowel disease. *J Crohns Colitis*. 2014;8(12):1661-1667. doi:10.1016/j.crohns.2014.07.007
53. Desai RJ, Pawar A, Khosrow-Khavar F, Weinblatt ME, Kim SC. Risk of venous thromboembolism associated with tofacitinib in patients with rheumatoid arthritis: a population-based cohort study. *Rheumatology (Oxford)*. 2021;61(1):121-130. doi:10.1093/rheumatology/keab294
54. Ma C, Panaccione R, Xiao Y, Khandelwal Y, Murthy SK, Wong ECL, et al. REMIT-UC: real-world effectiveness and safety of tofacitinib for moderate-to-severely active ulcerative colitis: a Canadian IBD Research Consortium Multicenter National Cohort Study. *Am J Gastroenterol*. 2023;118(5):861-871. doi:10.14309/ajg.0000000000002129
55. Lopetuso LR, Mocci G, Marzo M, D'Aversa F, Rapaccini GL,

- Guidi L, et al. Harmful effects and potential benefits of anti-tumor necrosis factor (TNF)- α on the liver. *Int J Mol Sci*. 2018;19(8). doi:10.3390/ijms19082199
56. Magri S, Chessa L, Demurtas M, Cabras F, Mocci G. Review article: safety of new biologic agents for inflammatory bowel disease in the liver. *Eur J Gastroenterol Hepatol*. 2021;33(5):623-630. doi:10.1097/meg.0000000000002076
 57. Kunchok A, Aksamit AJ, Jr., Davis JM, 3rd, Kantarci OH, Keegan BM, Pittcock SJ, et al. Association between tumor necrosis factor inhibitor exposure and inflammatory central nervous system events. *JAMA Neurol*. 2020;77(8):937-946. doi:10.1001/jamaneurol.2020.1162
 58. Kumar D, Bouldin TW, Berger RG. A case of progressive multifocal leukoencephalopathy in a patient treated with infliximab. *Arthritis Rheum*. 2010;62(11):3191-3195. doi:10.1002/art.27687
 59. Chow S, Patnana S, Gupta NK. Posterior Reversible encephalopathy syndrome in a patient with Crohn's disease on infliximab. *J Clin Gastroenterol*. 2016;50(8):687. doi:10.1097/mcg.0000000000000557
 60. Çimen Güneş E, Çolak S, Tekgöz E, Çınar M, Yılmaz S. Golimumab-induced posterior reversible encephalopathy syndrome (PRES): a case-based review. *Clin Rheumatol*. 2023;42(12):3407-3410. doi:10.1007/s10067-023-06771-w
 61. Jordan A, Kinnucan J. Ustekinumab-associated posterior reversible encephalopathy syndrome in a patient with Crohn's disease. *ACG Case Rep J*. 2022;9(10):e00867. doi:10.14309/crj.0000000000000867
 62. Mishra A, Seril DN. Posterior reversible encephalopathy syndrome following ustekinumab induction for Crohn's disease. *Case Rep Gastroenterol*. 2018;12(2):521-527. doi:10.1159/000492462
 63. Askling J, Fahrback K, Nordstrom B, Ross S, Schmid CH, Symmons D. Cancer risk with tumor necrosis factor alpha (TNF) inhibitors: meta-analysis of randomized controlled trials of adalimumab, etanercept, and infliximab using patient level data. *Pharmacoepidemiol Drug Saf*. 2011;20(2):119-130. doi:10.1002/pds.2046
 64. Axelrad J, Bernheim O, Colombel JF, Malerba S, Ananthakrishnan A, Yajnik V, et al. Risk of new or recurrent cancer in patients with inflammatory bowel disease and previous cancer exposed to immunosuppressive and anti-tumor necrosis factor agents. *Clin Gastroenterol Hepatol*. 2016;14(1):58-64. doi:10.1016/j.cgh.2015.07.037
 65. Nyboe Andersen N, Pasternak B, Basit S, Andersson M, Svanström H, Caspersen S, et al. Association between tumor necrosis factor- α antagonists and risk of cancer in patients with inflammatory bowel disease. *Jama*. 2014;311(23):2406-2413. doi:10.1001/jama.2014.5613
 66. Cohen JA, Khatri B, Barkhof F, Comi G, Hartung HP, Montalban X, et al. Long-term (up to 4.5 years) treatment with fingolimod in multiple sclerosis: results from the extension of the randomised TRANSFORMS study. *J Neurol Neurosurg Psychiatry*. 2016;87(5):468-475. doi:10.1136/jnnp-2015-310597
 67. Sandborn WJ, D'Haens GR, Sands BE, Panaccione R, Ng SC, Lawendy N, et al. Tofacitinib for the treatment of ulcerative colitis: an integrated summary of up to 7.8 years of safety data from the Global Clinical Programme. *J Crohns Colitis*. 2023;17(3):338-351. doi:10.1093/ecco-jcc/jjac141
 68. Curtis JR, Yamaoka K, Chen YH, Bhatt DL, Gunay LM, Sugiyama N, et al. Malignancy risk with tofacitinib versus TNF inhibitors in rheumatoid arthritis: results from the open-label, randomised controlled ORAL Surveillance trial. *Ann Rheum Dis*. 2023;82(3):331-343. doi:10.1136/ard-2022-222543